

# Colonoscopy Prep – Moviprep Split Dose Prep

Please review this entire document! These instructions are your physician’s specific instructions. Follow all steps carefully to ensure successful prep and procedure. If you are or may be pregnant, please discuss the risks and benefits of this procedure with your Provider. **Questions? Call 317.214.5468.**

Please do your best to keep your scheduled appointment time. If you must cancel or Reschedule, please call us at 317.214.5468 at least **SEVEN DAY’S NOTICE.**

<p><b>10 Days</b> before your procedure</p>	<p><b>Review and start following all MEDICATION GUIDELINES <a href="#">here.</a></b></p> <ul style="list-style-type: none"> <li>• Consult with your prescribed provider before discontinuing any blood thinners/ anticoagulants and/or diabetic medications.</li> </ul>
<p><b>1 Day</b> before your procedure</p>	<p><b>Start following CLEAR LIQUID DIET included in this document.</b></p> <ul style="list-style-type: none"> <li>• NO SOLID FOODS, CLEAR LIQUIDS ONLY</li> <li>• Drink plenty of clear liquids throughout the day to avoid dehydration.</li> </ul> <p>Each <b>MoviPrep</b> kit contains a disposable container, 2 pouches labeled A, and 2 pouches labeled B. You must complete the entire prep to ensure the most effective cleansing.</p> <p><b>AT 5PM:</b></p> <p><b>Step 1:</b> Mix the first dose - Empty 1 pouch A and 1 pouch B into the disposable container Add lukewarm drinking water to the top line of the container. Mix to dissolve <b>If preferred, mix the solution ahead of time and refrigerate prior to drinking.</b> The reconstituted solution <b>should be used within 24 hours.</b></p> <p><b>Step 2:</b> The MoviPrep container is divided into 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is consumed Drink 16 oz of the clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep. Take three Simethicone (GAS-X) 125mg tablets available over the counter.</p> <p><b>IMPORTANT:</b> The 2nd dose will be <b>6 hours prior to appointment arrival time</b> on the day of procedure.</p>
<p><b>The Day</b> of your procedure</p>	<p><b>6 HOURS PRIOR TO APPOINTMENT ARRIVAL TIME:</b> Drink second dose of MoviPrep by repeating Steps 1 and 2. Continue following CLEAR LIQUID DIET until 4 hours prior to appointment arrival time.</p> <p><b>4 HOURS PRIOR TO APPOINTMENT ARRIVAL TIME:</b></p> <ul style="list-style-type: none"> <li>• STOP DRINKING, NOTHING BY MOUTH</li> <li>• No eating, drinking, smoking, or vaping. No gum or hard candy.</li> <li>• Take any approved medication with a small sip of water at least 4 hours before the procedure.</li> <li>• You may brush your teeth, but do not swallow.</li> </ul>

- You are ready for the colonoscopy if you follow all instructions, and your stool is clear enough to see the bottom of the toilet bowl after a bowel movement.



# Colonoscopy Prep –

## Moviprep Split Prep

### BEFORE THE PROCEDURE

- If you are on blood thinners/anticoagulants, please contact your prescribed physician before discontinuing your medication. Medication guidelines are included [here](#).
- Colonoscopy prep will likely require the purchase of some over-the-counter medications and drinks allowed on the clear liquid diet. Shopping list allowed foods and foods not allowed are included in this document.
- If you are being sedated, please plan to have a friend/family member adult driver (age 18+) for after your procedure.

### THE PREP

- Be sure to have access to a restroom. Individual response time to prep may vary. You may have loose stools.
- You may be uncomfortable/bloated and feel chilled until you start having bowel movements (2-4 hours). Drinking warm, clear liquids such as tea, warm apple juice, or hot water can help.
- Alcohol-free baby wipes may help ease skin irritation.
- Use over-the-counter hemorrhoid pads if necessary.

### THE APPOINTMENT

- Bring to the appointment:
  - List of all current medications (including allergy or over the counter)
  - Inhalers
  - Photo ID and up-to-date insurance information
  - Leave valuables (jewelry, smart watches, cell phones) at home or with your driver
- Arrive 1 hour prior to scheduled procedure.
- You **MUST** have a friend/family member adult driver (age 18+) remain in the building throughout the entire procedure and available to drive you home after your procedure.
- Taxi/cab, Uber/Lyft, bus and medical transport are **NOT** acceptable rides for home unless accompanied by an adult (age18+) friend/family member.

## **AFTER THE PROCEDURE**

Because of the sedation given during the procedure, you may experience sleepiness, dizziness, forgetfulness, or lightheadedness. Your judgement and reflexes may be impaired. These are normal reactions to the medication that can last for the remainder of the day.

### **FOR THE REMAINDER OF YOUR PROCEDURE DAY**

- **DO NOT** drive a vehicle or operate complex machinery
- **DO NOT** perform any strenuous activity
- **DO NOT** drink alcohol
- **DO NOT** smoke unsupervised
- **DO NOT** make any legal decisions
- Additional information will be included on your discharge instructions

### **SHOPPING LIST**

#### **Required Items**

- Allowed beverages for clear liquid diet
- Moviprep Kit
- Three Simethicone (GAS-X) 125mg tablets available over the counter.



#### **Optional Items**

- Hemorrhoidal pads
- Alcohol-free baby wipes

## **CLEAR LIQUID DIET**

### **NO SOLID FOODS. CLEAR LIQUIDS ONLY.**

Be sure to drink plenty of clear liquids with calories on clear liquid diet days

### **APPROVED CLEAR LIQUIDS**

- Sports drinks/Gatorade®/VitaminWater® No red or purple
- Fruit juice (no pulp) Apple, white grape, white cranberry; No red or purple
- Soda & carbonated beverages; No red or purple
- Clear broth/bouillon Chicken, vegetable, beef flavors
- Gelatin/Jell-O® No red or purple
- Ice pop/Popsicles® No red or purple
- Gummy bears; No red or purple
- Black coffee & tea No liquid or powdered creamer/milk (dairy/soy/nut/oat);  
Sugar/sweeteners OK
- Water Plain, flavored, carbonated, No red or purple

### **IF YOU ARE DIABETIC, AIM FOR 45 GRAMS OF CARBOHYDRATES PER MEAL AND 15-30 GRAMS PER SNACK**

- **Clear liquids with about 15 grams of carbohydrates**
  - 4oz apple juice, 8oz sports drink, ½ cup gelatin, 2 popsicles/ice pops
  - Clear liquids with zero carbohydrates
  - Black coffee, tea (unsweetened or diet), clear diet soda, seltzer, flavored water, fat-free broth, bouillon or consommé

**⊘ NO RED ⊘ NO PURPLE ⊘ NO DAIRY ⊘ NO PULP**